



Pavella Creative **Design**

Biohungaricum logo és arculattervezés



BIOHUNGARICUM
From Genuine Source





Egészségügyi információs szolgáltatás
Életet menthet!



MNG logo és arculattervezés

MAGYAR
NEMZETI
galéria



MNG
MAGYAR
NEMZETI
GALÉRIA





Budapest logo és arculattervezés

BUDAPEST

GET BALANCED



Telki község arculatfrissítés






Paprika Gourmet logo és
arculattervezés



Paprika Gourmet
Hungarian Delicatessen







Angiocarb logo és
arcuartervezés




angiocarb®

Menta magazin logo és
arculattervezés




Menta magazin online kiadványtervezés

Mediso arculat tervezés



nanoScan® PET
Preclinical Imaging Systems



nanoScan® PET/MRI
nanoScan® PET/CT



IMAGING FOR SCIENCE

nanoScan® PC PET/CT
in vivo molecular and preclinical imager



The first sub-half mm³ PET volumetric resolution PET/CT with exceptional image quantification - accuracy over 97 %

Preclinical Line



IMAGING FOR SCIENCE

MEDISO GROUP OF COMPANIES




The most flexible clinical and preclinical imaging portfolio



IMAGING FOR LIFE

nanoScan® Family
the next generation four modality preclinical imaging platform
SPECT/MRI, PET/MRI, SPECT/CT, PET/CT, SPECT/CT/PET



nanoScan® - leading performance in all modalities

- breakthrough 13,000 cps/MBq SPECT sensitivity
- 300 µm SPECT and 700 µm PET resolution
- quantification accuracy over 97% for SPECT & PET
- Fully integrated SPECT/MRI and PET/MRI systems



IMAGING FOR LIFE

LEUCO-SCINT kit

Separation and labelling protocol (Aseptic conditions should be kept throughout the separation and labelling process)

1. Draw 2 ml of ACD solution and 3 ml of Fluorescein into each of the four 20 ml sterile plastic syringes.
2. Withdraw 15 ml patient's blood into each syringe and mix gently by inversion. Note in case of infants the labelling can be managed with 2-3 ml.
3. Leave to stand the syringes for 30-40 mins to allow red blood cells to sediment at room temperature.
4. When the red cells are sedimented to about half of the original volume of the blood, carefully draw up the leucocytes platelet rich plasma (LPP) into sterile tubes and centrifuge 10 mins at 150 g.
5. Remove and retain all the supernatant (platelet rich plasma, PRP). Pool the pellets of mixed leucocytes in one sterile tube. Centrifuge again for 10 mins at 150 g. Remove the remaining supernatant from the pellet of mixed leucocytes, leaving the pellet almost to dry. Note 1: During centrifugation phase start labelling a Leuco-Scint kit vial with 10-90s. Note 2: only 15 ml PRP is needed to procedure described in step 7.
6. Reconstitute one vial of LEUCO-SCINT with 1.5 ml eluate of ^{99m}Tc -generator containing 100-750 MBq ^{99m}Tc -pertechnetate. Shake for 10 mins to dissolve the HM-PAO and add exactly 1 ml immediately ^{99m}Tc -HM-PAO to a tube of mixed leucocytes (the activity of ^{99m}Tc -HM-PAO is approx. 400-500 MBq). Mix gently and incubate the cells for 10 mins at room temperature.
7. While incubating the leucocytes, centrifuge the PRP (step 5) for 5 mins at 2000 g to produce cell-free plasma (CFP).
8. Add 3.5 ml CFP (obtained in step 7) to the labelled cell suspension and gently mix. Centrifuge at 150g for 10 mins.
9. Transfer all supernatant to a tube.
10. Add 3.5 ml of CFP to the pellet of leucocytes, gently swirl to mix.
11. Measure the radioactivity and calculate the cell labelling efficiency (this activity of labelled leucocytes activity of labelled leucocytes ÷ activity of supernatant × 100).
12. The labelled leucocytes should be re-injected without delay. Imaging at 30 mins, 2 hrs and 24 hrs after administration.



Medi Radiopharma arcuat tervezés

MEDI-RADIOPHARMA

^{99m}Tc jelzésre alkalmas in vivo kitek egyre bővülő széles választéka

CARDIO-SPECT 0,5 mg
(Európai regisztráció)

Stabilizált BRAIN-SPECT kit

ALBUMON kit (^{99m}Tc -HSA)

MEDI-RADIOPHARMA Co., Ltd
2030 Érd, Szamos utca 10-12.
Telefon: 06-23/521-261
Fax: 06-23/521-260
e-mail: order@mediradiopharma.hu
www.mediradiopharma.com

MEDI-RADIOPHARMA

Minőségellenőrző készletek
a GMP követelményeknek megfelelő aszeptikus laboratóriumi munkákhoz

MediCheck Q.C. kit

Medi-Media Fill kit

MEDI-RADIOPHARMA Co., Ltd
2030 Érd, Szamos utca 10-12.
Telefon: 06-23/521-261
Fax: 06-23/521-260
e-mail: order@mediradiopharma.hu
www.mediradiopharma.com

MEDI-RADIOPHARMA

Sugárvédelmi eszközök és berendezések
radioaktív gyógyszer előállításához szükséges eszközök és kellékek

GammaFinder II.
vezeték nélküli detektor intraoperatív felhasználásra

Cégünk a kizárólagos magyarországi forgalmazó

MEDI-RADIOPHARMA Co., Ltd
2030 Érd, Szamos utca 10-12.
Telefon: 06-23/521-261
Fax: 06-23/521-260
e-mail: order@mediradiopharma.hu
www.mediradiopharma.com

MEDI-RADIOPHARMA

20 évesek lettünk!
a MEDI-RADIOPHARMA 20 éve az Önök szolgálatában az Önök támogatásával és együttműködésével.

MEDI-RADIOPHARMA Co., Ltd
2030 Érd, Szamos utca 10-12.
Telefon: 06-23/521-261
Fax: 06-23/521-260
e-mail: order@mediradiopharma.hu
www.mediradiopharma.com

DICSŐSÉG AZ ULTRÁKNAK

GLORY TO THE ULTRAS



ULTRABALATON

A FRISSÍTÉS TOVÁBBRA IS PRIMAVERA ÉS POWERBAR


REFRESHMENTS ARE STILL
PRIMAVERA AND POWERBAR



ULTRABALATON

CÉGES ÉS EGYETEMI CSAPATOK


COMPANY AND UNIVERSITY TEAMS



ULTRABALATON

AZ IRÁNY VÁLTOZATLAN

THE SAME DIRECTION



ULTRABALATON



ULTRABALATON

HÁROM FŐS CSAPATOK

THREE-PERSON TEAMS



ULTRABALATON

Ggggú babaságok – logótervezés



Egyéb logók



Egyéb logók






szerkesztette: Sakonyi Péter
ára: 980 forint



Egyéb kiadványok



Jelen dokumentum a Pavella Bt. szellemi terméke.
A dokumentum harmadik félnek sem részben sem egészben
nem adható ki és nem publikálható.

